

## An exact multiplicity result for a class of symmetric problems

Philip Korman

(Communicated by Luís Sanchez)

**Abstract.** We consider positive solutions of a class of semilinear problems

$$u'' + \lambda a(x)f(u) = 0, \quad -1 < x < 1, \quad u(-1) = u(1) = 0,$$

with even and positive  $a(x)$ , depending on a positive parameter  $\lambda$ . In case  $f(u)$  is convex, an exact multiplicity result was given in P. Korman, Y. Li and T. Ouyang [6]; see also P. Korman [4] for the details. It was observed by P. Korman and J. Shi [7] that convexity requirement can be relaxed for large  $u$  (see also [5]). We show that convexity requirement can also be relaxed for small  $u$ .

**Mathematics Subject Classification (2000).** 34B15.

**Keywords.** Exact multiplicity of solutions.

We consider positive solutions of a class of non-autonomous problems

$$u'' + \lambda a(x)f(u) = 0, \quad -1 < x < 1, \quad u(-1) = u(1) = 0, \quad (1)$$

with even and positive  $a(x)$ , depending on a positive parameter  $\lambda$ . We assume that the function  $a(x) \in C^1(-1, 1) \cap C[-1, 1]$  satisfies

$$a(x) > 0, \quad a(-x) = a(x), \quad a'(x) < 0 \quad \text{for } x \in (0, 1), \quad (2)$$

while  $f(u) \in C^2(\bar{R}_+)$  satisfies

$$f(u) > 0 \quad \text{for } u \geq 0. \quad (3)$$

We study *exactly* how many solutions the problem (1) has, and how these solutions are connected when  $\lambda$  is varied.

It follows from B. Gidas, W.-M. Ni and L. Nirenberg [3] that under these conditions any positive solution of (1) is an even function, with  $u'(x) < 0$  for  $x > 0$

(i.e.,  $u(0)$  gives the maximum value of the solution). It is also known that in this case the problem (1) has properties similar to those of autonomous problems; see a recent review paper [4]. In particular, any non-trivial solution of the corresponding linearized problem

$$w'' + \lambda a(x)f'(u)w = 0, \quad -1 < x < 1, \quad w(-1) = w(1) = 0, \quad (4)$$

is an even function of one sign, so that we may assume that  $w(x) > 0$  on  $(-1, 1)$ . It is also known that as one varies  $\lambda$ , solutions of (1) lie on smooth solution curves, which admit only simple turns at singular solutions (i.e., solutions  $(\lambda, u(x))$  of (1), at which the problem (4) admits non-trivial solutions).

We now make a further assumption on  $f(u)$ . We assume there is an  $\gamma > 0$ , so that

$$f'(u) < 0 \quad \text{for } 0 < u < \gamma, \quad f'(u) > 0 \quad \text{and} \quad f''(u) > 0 \quad \text{for } u > \gamma. \quad (5)$$

(Observe that we do not restrict convexity on the interval  $(0, \gamma)$ .)

**Lemma 1.** *Under the assumption (5), for any singular solution  $u(x)$  and the corresponding non-trivial solution of (4) one has*

$$p(x) \equiv 3u'(x)w'(x) - u''(x)w(x) > 0 \quad \text{for all } x \in [0, 1].$$

*Proof.* It follows by maximum principle that (4) cannot have a non-trivial solution in the region where  $f'(u) < 0$ , i.e.  $u(0) > \gamma$  at any singular solution. Since  $w(x) > 0$ , we see from the equation (4) that  $w(x)$  changes concavity exactly once on  $(0, 1)$ , being concave for small  $x$  and convex near  $x = 1$ . It follows that  $w(x)$  is a non-increasing function, i.e.,  $w'(x) \leq 0$  for all  $x \in (0, 1)$ . But then  $p(x) = 3u'(x)w'(x) + \lambda a(x)f(u(x))w(x) > 0$ .  $\square$

**Theorem 1.** *For the problem (1) assume that the conditions (2), (3) and (5) hold. Then the problem (1) has at most two positive solutions, and moreover all solutions lie on a unique smooth solution curve, starting at  $(\lambda = 0, u = 0)$ . This curve makes at most one turn, and it tends to infinity at some  $\bar{\lambda} \geq 0$ . If, in addition, we assume that*

$$\lim_{u \rightarrow \infty} \frac{f(u)}{u} = \infty, \quad (6)$$

*then this curve of solutions makes exactly one turn at some  $\lambda = \lambda_0$ , and it tends to infinity as  $\lambda \downarrow 0$ , so that for  $0 < \lambda < \lambda_0$  the problem (1) has exactly two strictly ordered positive solutions, it has exactly one positive solution at  $\lambda = \lambda_0$ , and none for  $\lambda > \lambda_0$ .*

*Proof.* By the implicit function theorem there is a curve of positive solutions, starting at  $(\lambda = 0, u = 0)$ . We can continue this curve for increasing  $\lambda$  by using

either the implicit function theorem or the Crandall–Rabinowitz bifurcation theorem [2]; see [4] for the details. Since  $f(u)$  is increasing and convex for large  $u$ , it follows that it grows at least linearly, and then the Sturm comparison theorem implies that this curve cannot extend for all  $\lambda > 0$ , so that it will either go to infinity at a finite  $\bar{\lambda}$ , or it will turn left eventually. If condition (6) holds, then the second possibility must be true, again by the Sturm comparison theorem.

The key is to show that a turn to the left in the  $(\lambda, u(0))$  plane occurs at any turning point. This will imply that there is only one turning point, and the theorem will follow.

Let  $u(x)$  be a singular solution of (1). We denote by  $\alpha = u(0)$  its maximal value. By maximum principle  $\alpha > \gamma$ . It then follows by our assumption (5) that

$$f'(u(x)) - f'(\alpha) < 0 \quad \text{for all } x \in (0, 1). \quad (7)$$

The direction of turn at any critical solution of (1) is governed by the sign of the integral  $I \equiv \int_0^1 a(x)f''(u)w^3 dx$ . We show next that this integral is positive, which implies that a turn to the left must occur at any singular solution; see, e.g., [4]. Indeed

$$\begin{aligned} I &= \int_0^1 \frac{d}{dx} [f'(u) - f'(\alpha)] \frac{a(x)w^3(x)}{u'(x)} dx \\ &= [f'(u) - f'(\alpha)] \frac{aw^3}{u'} \Big|_0^1 - \int_0^1 [f'(u) - f'(\alpha)] \frac{a'u'w^3 + aw^2p}{u'^2} dx. \end{aligned} \quad (8)$$

The first term on the right vanishes, since

$$\lim_{x \rightarrow 0} \frac{f'(u) - f'(\alpha)}{u'} = \lim_{x \rightarrow 0} f''(u)u'u'' = 0, \quad (9)$$

while the second term is positive by (7) and Lemma 1. (Observe that the integrand in the second term is bounded, as seen by a computation similar to (9).)  $\square$

We remark that our result might be new even for constant  $a(x)$ , although in that case it can probably be obtained by the time-map method; see, e.g., I. Addou and S.-H. Wang [1].

## References

- [1] I. Addou and S.-H. Wang, Exact multiplicity results for a  $p$ -Laplacian positone problem with concave-convex-concave nonlinearities. *Electron. J. Differential Equations* **2004** (2004), no. 72, 1–24. [Zbl 1057.34008](#) [MR 2057659](#)

- [2] M. G. Crandall and P. H. Rabinowitz, Bifurcation, perturbation of simple eigenvalues and linearized stability. *Arch. Rational Mech. Anal.* **52** (1973), 161–180. [Zbl 0275.47044](#) [MR 0341212](#)
- [3] B. Gidas, W. M. Ni, and L. Nirenberg, Symmetry and related properties via the maximum principle. *Comm. Math. Phys.* **68** (1979), 209–243. [Zbl 0425.35020](#) [MR 544879](#)
- [4] P. Korman, Global solution branches and exact multiplicity of solutions for two point boundary value problems. In *Handbook of differential equations: Ordinary differential equations*, vol. 3, North Holland, Amsterdam 2006, 547–606.
- [5] P. Korman, Stability and instability of solutions of semilinear problems. *Appl. Anal.* **86** (2007), 135–147. [Zbl 1127.34008](#) [MR 2297310](#)
- [6] P. Korman, Y. Li, and T. Ouyang, Exact multiplicity results for boundary value problems with nonlinearities generalising cubic. *Proc. Roy. Soc. Edinburgh Sect. A* **126** (1996), 599–616. [Zbl 0855.34022](#) [MR 1396280](#)
- [7] P. Korman and J. Shi, Instability and exact multiplicity of solutions of semilinear equations. In *Proceedings of Nonlinear Differential Equations* (Coral Gables, FL, 1999), Electron. J. Differ. Equ. Conf. 5, Southwest Texas State Univ., San Marcos, TX 2000, 311–322. [Zbl 0970.34015](#) [MR 1799061](#)
- [8] R. Schaaf and K. Schmitt, A class of nonlinear Sturm-Liouville problems with infinitely many solutions. *Trans. Amer. Math. Soc.* **306** (1988), 853–859. [Zbl 0657.34021](#) [MR 933322](#)

Received March 6, 2007; received May 9, 2007

P. Korman, Department of Mathematical Sciences, University of Cincinnati, Cincinnati, OH 45221-0025, U.S.A.

E-mail: [kormanp@math.uc.edu](mailto:kormanp@math.uc.edu)