

Matrix Transformations

Introduction to Linear Transformations

Linear Algebra
MATH 2076



Another look at the matrix product $A\vec{x}$

Let \vec{a}_j be the j^{th} column of some $m \times n$ matrix A , and \vec{x} be in \mathbb{R}^n , so

$$A = \begin{bmatrix} a_{11} & a_{12} & \cdots & a_{1n} \\ a_{21} & a_{22} & \cdots & a_{2n} \\ \vdots & \vdots & \vdots & \vdots \\ a_{m1} & a_{m2} & \cdots & a_{mn} \end{bmatrix}, \quad \vec{a}_j = \begin{bmatrix} a_{1j} \\ a_{2j} \\ \vdots \\ a_{mj} \end{bmatrix}, \quad \text{and} \quad \vec{x} = \begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{bmatrix}.$$

Recall that the product $A\vec{x}$ is defined to be the LC

$$x_1\vec{a}_1 + x_2\vec{a}_2 + \cdots + x_n\vec{a}_n.$$

Since each \vec{a}_j is a vector in \mathbb{R}^m , so is $A\vec{x}$. Thus we can define a vector function by the rule $\vec{y} = A\vec{x}$. Here

$$\begin{bmatrix} y_1 \\ y_2 \\ \vdots \\ y_m \end{bmatrix} = \vec{y} = A\vec{x} = x_1\vec{a}_1 + x_2\vec{a}_2 + \cdots + x_n\vec{a}_n.$$

The matrix transformation $\vec{y} = A\vec{x}$

This defines a *function* from \mathbb{R}^n to \mathbb{R}^m ; the input variable \vec{x} comes from \mathbb{R}^n , it gets multiplied by the matrix A via the formula

$$x_1\vec{a}_1 + x_2\vec{a}_2 + \cdots + x_n\vec{a}_n,$$

and we call the resulting output \vec{y} . That is,

$$\begin{bmatrix} y_1 \\ y_2 \\ \vdots \\ y_m \end{bmatrix} = \vec{y} = A\vec{x} = x_1\vec{a}_1 + x_2\vec{a}_2 + \cdots + x_n\vec{a}_n.$$

In linear algebra, functions are usually called *transformations*.

We write $\mathbb{R}^n \xrightarrow{T} \mathbb{R}^m$ to indicate that T is a transformation from \mathbb{R}^n to \mathbb{R}^m , meaning that the input variable \vec{x} comes from \mathbb{R}^n and the resulting output $\vec{y} = T(\vec{x})$ is a vector in \mathbb{R}^m .

A transformation (aka function) $\mathbb{R}^n \xrightarrow{T} \mathbb{R}^m$

Here \mathbb{R}^n is the *domain* of T — where the input variables \vec{x} live, and \mathbb{R}^m is the *codomain* of T — where the resulting output $\vec{y} = T(\vec{x})$ lives.

For each \vec{x} in \mathbb{R}^n , $\vec{y} = T(\vec{x})$ is called the *T -image* of \vec{x} . If \mathbb{S} is a bunch of vectors in \mathbb{R}^n (i.e., $\mathbb{S} \subset \mathbb{R}^n$), then

$$T(\mathbb{S}) = \{\text{all images } T(\vec{x}) \text{ where } \vec{x} \text{ is in } \mathbb{S}\}$$

is called the *T -image* of \mathbb{S} .

The *range* of T is $\mathcal{R}ng(T) = T(\mathbb{R}^n)$, the set of all images $T(\vec{x})$; \vec{b} is in $\mathcal{R}ng(T)$ if and only if $\vec{b} = T(\vec{x})$ for some \vec{x} . Evidently, $\mathcal{R}ng(T)$ is a subset of the codomain of T ; $\mathcal{R}ng(T) \subset \mathbb{R}^m$.

An important question is to know which vectors \vec{b} are in the range of T . That is: Given \vec{b} , when can we find an \vec{x} in \mathbb{R}^n with $T(\vec{x}) = \vec{b}$?

A matrix transformation $\mathbb{R}^n \xrightarrow{T} \mathbb{R}^m$

Here we assume that $\mathbb{R}^n \xrightarrow{T} \mathbb{R}^m$ is given by the rule $T(\vec{x}) = A\vec{x}$ for some $m \times n$ matrix A . So, for each \vec{x} in \mathbb{R}^n ,

$$T(\vec{x}) = A\vec{x} = x_1\vec{a}_1 + x_2\vec{a}_2 + \cdots + x_n\vec{a}_n$$

where \vec{a}_j is the j^{th} column of A . Thus each image $T(\vec{x})$ is a **LC** of the columns of A .

Therefore, the *range* of T , which is the set of all images $T(\vec{x})$, is the set of all linear combinations of the columns of A ; i.e., the range of T is the *span* of the columns of A .

Look at: Given \vec{b} , when can we find an \vec{x} in \mathbb{R}^n with $T(\vec{x}) = \vec{b}$? Here we are just asking whether or not we can solve $A\vec{x} = \vec{b}$.

The *range* of T is exactly all rhs vectors \vec{b} such that $A\vec{x} = \vec{b}$ has a solution.

The range of $\mathbb{R}^n \xrightarrow{T} \mathbb{R}^m$

Assume $\mathbb{R}^n \xrightarrow{T} \mathbb{R}^m$ is given by $T(\vec{x}) = A\vec{x}$ for some $m \times n$ matrix A .

The *range* of T is $\mathcal{Rng}(T) = T(\mathbb{R}^n)$ (the T -image of \mathbb{R}^n).

There are many equivalent ways to view the range of T !

- $\mathcal{Rng}(T)$ is all images $T(\vec{x})$ for \vec{x} in \mathbb{R}^n
- $\mathcal{Rng}(T)$ is all \vec{b} in \mathbb{R}^m such that $\vec{b} = T(\vec{x})$ for some \vec{x} in \mathbb{R}^n
- $\mathcal{Rng}(T)$ is all \vec{b} in \mathbb{R}^m such that $A\vec{x} = \vec{b}$ has a solution (is consistent)
- $\mathcal{Rng}(T)$ is the *span* of the columns of A