

Adventures in the $\Upsilon(10860)$ Neighborhood at Belle



- Belle/KEKB: B-factory
[$\Upsilon(4S)$ Resonance, B meson]
- $\Upsilon(10860)$ Region
 - physics interest
 - spectroscopic anomalies
 - future



Kay Kinoshita
University of Cincinnati
Belle Collaboration






Belle collaboration




International, ~400-600 collaborators

Belle hardware



- Charged tracking/vertexing
 - SVD:
 - (-7/03) 3-layer DSSD Si μstrip (152M B pairs)
 - (8/03-) 4-layer (550+M BB)
 - CDC: 50 layers (He-ethane)
- Hadron identification
 - CDC: dE/dx
 - TOF: time-of-flight
 - ACC: Threshold Cerenkov (aerogel)
- Electron/photon
 - ECL: CsI calorimeter
- Muon/ K_L
 - KLM: Resistive plate counter/iron



4/07- COPPER pipelined DAQ system


... the main thrust

$$e^+ e^- \rightarrow \Upsilon(4S) \rightarrow B\bar{B}$$



Virginia Tech, April 1, 2015

K. Kinoshita



“B-factory”

Primary goal: discover CP violation in weak decays of B meson

DONE! (2001)

Data (1999-2010)



> 1 ab⁻¹ recorded by Belle

$\int L dt$ since 6/1999

- $\Upsilon(4S)$
 711 fb⁻¹
- sub- $\Upsilon(4S)$ continuum
 ~100 fb⁻¹
- “ $\Upsilon(5S)$ ”
 ~121 fb⁻¹
- $\Upsilon(3S)$, $\Upsilon(2S)$, $\Upsilon(1S)$
 ~34 fb⁻¹
- “ $\Upsilon(5S)$ ”+ scan
 ~31 fb⁻¹



- B pairs (7.7×10^8 events)
- charm (1.1×10^9 events)
- tau ($\sim 8 \times 10^8$ events)
- 2-photon events
- B_s ($\sim 7 \times 10^6$ events)
- $\Upsilon(5S)$ ($\sim 4 \times 10^7$ events)



Belle physics

434 articles published/submitted

http://belle.kek.jp/bdocs/b_journal.html


Highlights

- CP violation in B decay
- Constraints on CKM; precision $\sin 2\varphi_1$, $|V_{cb}|$, $|V_{ub}|$
- overconstraints on CKM; limits/hints on New Physics
- evidence for D^0 mixing
- new charmonium-like states $Z(4430)$, $\Upsilon(4660)$, $\Upsilon(4008)$,
 $X(4160)$, $\Upsilon(3940)$, $X(3872)$
- new bottomonia, bottomonium-like $Z_b(10610)$, $Z_b(10650)$
- Kobayashi & Maskawa 2008 Nobel

Future: Super KEKB / Belle II

- to start ~ 2016

The Upsilon Neighborhood



At/above the $\Upsilon(10860)$ [" $\Upsilon(5S)$ "]: B_s , bottomonium physics

- ✧ B-factory detector: high luminosity, established detector, $\Upsilon(4S)$ data for comparison; CLEAN events, energy definition, γ detection; high trigger efficiency
- ✧ on resonance - # events measured directly \rightarrow absolute BF's

Belle data above $\Upsilon(4S)$: history

2005: 3-day “engineering” run

- basic $\Upsilon(10860)$, $B_s^{(*)}$ properties,
- test KEKB at $\Upsilon(10860)$
- 1.86 fb^{-1} at peak (10869 MeV)

A. Drutskoy et al., PRL 98, 052001 (2007)

A. Drutskoy et al., PRD 76, 012002 (2007)

June 2006: 20-day run

- + 21.7 fb^{-1} on resonance

K.F. Chen et al., PRL 100, 112001 (2008)

J. Wicht et al., PRL 100, 121801 (2008)

R. Louvot et al., PRL 102, 021801 (2009)


A. Drutskoy et al., PRD 81, 112003(R)(2010)

R. Louvot et al., PRL 104, 231801 (2010)

C.-C. Peng et al., PRD 82, 072007 (2010)

S. Esen et al., PRL 105, 201802 (2010)

J. Li et al., PRL 106, 121802 (2011)



Green: states including bottomonia
Black: B_s , other

2007: scan 6 pts

- + 7.9 fb^{-1} above resonance

K.F. Chen et al., PRD 82, 091106(R) (2010)

Belle data above $\Upsilon(4S)$



10/08-12/10: extended run

- $\sim 100 \text{ fb}^{-1}$ on resonance

I. Adachi et al, PRL 108, 032001 (2012)

A. Bondar et al, PRL 108, 122001 (2012)

Y. Sato et al, PRL 108, 171801 (2012)

J. Li et al, PRL 108, 181808 (2012)

R. Mizuk et al, PRL 109, 232002 (2012)

S. Esen et al, PRD 87, 031101(R) (2013)

C. Oswald et al, PRD 87, 072008 (2013)

E. Solovieva et al, PLB 726, 206 (2013)

P. Krokovny et al, PRD 88, 052016 (2013)

F. Thorne et al, PRD 88, 114006 (2013)


A. Garmash et al, arXiv:1403.0992 [accepted PRD]

X. He et al, PRL 113, 142001 (2014)


D. Dutta et al, PRD 91, 011101 (R) (2015)

- $\sim 30 \text{ fb}^{-1}$ scan $\Upsilon(4S) \rightarrow \Upsilon(11020)$

D. Santel et al, arXiv:1501.01137 [PRL]



Green: states including bottomonia
Black: B_s , other




$\Upsilon(10860)$ physics

Virginia Tech, April 1, 2015

K. Kinoshita

B_s decay in Standard Model

- similar to non-strange B spectator decay \rightarrow quark-hadron duality correspondence btw final particle ($D \leftrightarrow D_s$)
- dissimilarities
 - $\Delta\Gamma/\Gamma_{CP}/\Gamma = O(10\%)$
 - CP -asymmetry ~ 0
- In LHCb era: focus on final states w neutrals, absolute rates



spectroscopy

- $B_s^{(*)}$ mass
- $B_{(s)}^{(*)}(\pi)$ event fractions
- bottomonium, bottomonium-like states



Anomalies of the $\Upsilon(10860)$

(semi-historical progression)

Speculation [W.S. Hou, PRD 74, 017504 (2006)]



- 2005: charmonium-like particle at 4260 GeV found in

$$e^+ e^- \rightarrow \gamma_{ISR} \pi^+ \pi^- J/\psi \quad e^+ e^- \rightarrow \pi^+ \pi^- J/\psi$$

Babar PRL 95, 142001 (2005)

Belle PRD 77, 011105 (R) (2008)

CLEO PRD 74, 091104(R) (2006)

$$Y(4260) \rightarrow \pi^+ \pi^- J/\psi$$

+ many more! (now called X by PDG)

$$Y \rightarrow \pi^+ \pi^- \psi(2S)$$

- Does(do) analogous state(s) exist in Upsilon region, observable in $\Upsilon(5S)$ data?
- Might the $\Upsilon(10860)$ actually be a Υ state (Υ_b)?




Anomalies of the $\Upsilon(10860)$

✧ Search for

$$e^+ e^- \rightarrow \Upsilon(1S/2S/3S) \pi^+ \pi^-$$

PRL 100, 112001 (2008)

$\Upsilon(1S/2S/3S) \rightarrow \mu^+\mu^-$ 23.6 fb^{-1} Exclusive $\mu\mu\pi\pi$ events

4 modes seen $\Upsilon(10860) \rightarrow \Upsilon(nS)h^+h^-$

Process	$\sigma(\text{pb})$	$\mathcal{B}(\%)$	$\Gamma(\text{MeV})$
$\Upsilon(1S)\pi^+\pi^-$	$1.61 \pm 0.10 \pm 0.12$	$0.53 \pm 0.03 \pm 0.05$	$0.59 \pm 0.04 \pm 0.09$
$\Upsilon(2S)\pi^+\pi^-$	$2.35 \pm 0.19 \pm 0.32$	$0.78 \pm 0.06 \pm 0.11$	$0.85 \pm 0.07 \pm 0.16$
$\Upsilon(3S)\pi^+\pi^-$	$1.44^{+0.55}_{-0.45} \pm 0.19$	$0.48^{+0.18}_{-0.15} \pm 0.07$	$0.52^{+0.20}_{-0.17} \pm 0.10$
$\Upsilon(1S)K^+K^-$	$0.185^{+0.048}_{-0.041} \pm 0.028$	$0.061^{+0.016}_{-0.014} \pm 0.010$	$0.067^{+0.017}_{-0.015} \pm 0.013$

$\Upsilon(5S)$ expectation: width comparable to $\Upsilon(2S/3S/4S)$

Process	Γ_{total}	$\Gamma_{e^+e^-}$	$\Gamma_{\Upsilon(1S)\pi^+\pi^-}$
$\Upsilon(2S) \rightarrow \Upsilon(1S)\pi^+\pi^-$	0.032 MeV	0.612 keV	0.0060 MeV
$\Upsilon(3S) \rightarrow \Upsilon(1S)\pi^+\pi^-$	0.020 MeV	0.443 keV	0.0009 MeV
$\Upsilon(4S) \rightarrow \Upsilon(1S)\pi^+\pi^-$	20.5 MeV	0.272 keV	0.0019 MeV
$\Upsilon(10860) \rightarrow \Upsilon(1S)\pi^+\pi^-$	110 MeV	0.31 keV	0.59 MeV

larger by
 $> 10^2$

$\Upsilon(10860) = \Upsilon(5S)$, Υ_b , or something else?

→ 12/07: energy scan, for $e^+e^- \rightarrow \Upsilon(nS)h^+h^-$




Anomalies of the $\Upsilon(10860)$
resonant substructure in
 $e^+e^- \rightarrow \Upsilon(nS)\pi^+\pi^-$
 $e^+e^- \rightarrow h_b(mP)\pi^+\pi^-$
121 fb^{-1} at $\Upsilon(10860)$ peak

[PRL 108, 122001 (2012)]

$$\Upsilon(5S) \rightarrow \Upsilon(nS)\pi^\pm\pi^\mp$$

$$\Upsilon(5S) \rightarrow \Upsilon(nS)\pi^\pm\pi^\mp \quad \Upsilon(nS) \rightarrow \mu^+\mu^-$$




Full reconstruction,
Clean signal
→ Dalitz analysis

$$\Upsilon(5S) \rightarrow \Upsilon(nS)\pi^\pm\pi^\mp$$

18

Dalitz Plots 121.4 fb^{-1} $M^2(\Upsilon\pi)$ vs $M^2(\pi\pi)$



Fit function:

$$S(s_1, s_2) = |A_{Z_{b1}} + A_{Z_{b2}} + A_{\text{NR}} + A_{f_0(980)} + A_{f_2(1275)}|^2$$

$$s_i = M_{\pi_i}^2 \Upsilon \quad A_{Z_{bk}} = \frac{\sqrt{M_k \Gamma_k}}{M_k^2 - s_1 + i M_k \Gamma_k} + \frac{a_k e^{i \phi_k} \sqrt{M_k \Gamma_k}}{M_k^2 - s_2 + i M_k \Gamma_k}$$

$$A_{\text{NR}} = c_1 + c_2 M_{\pi\pi}^2$$

A. Voloshin, PRD74, 054022 (2006);
Prog. Part. Nucl. Phys. 61, 455 (2008)



2 new charged Bottomonium-like states

Final state	$\Upsilon(1S)\pi^+\pi^-$	$\Upsilon(2S)\pi^+\pi^-$	$\Upsilon(3S)\pi^+\pi^-$
$M(Z_b(10610))$	$10609 \pm 3 \pm 2$	$10616 \pm 2^{+3}_{-4}$	$10608 \pm 2^{+5}_{-2}$
$\Gamma(Z_b(10610))$	$22.9 \pm 7.3 \pm 2$	$21.1 \pm 4^{+2}_{-3}$	$12.2 \pm 1.7 \pm 4$
$M(Z_b(10650))$	$10660 \pm 6 \pm 2$	$10653 \pm 2 \pm 2$	$10652 \pm 2 \pm 2$
$\Gamma(Z_b(10650))$	$12 \pm 10 \pm 3$	$16.4 \pm 3.6^{+4}_{-6}$	$10.9 \pm 2.6^{+4}_{-2}$
Rel. amplitude	$0.59 \pm 0.19^{+0.09}_{-0.03}$	$0.91 \pm 0.11^{+0.04}_{-0.03}$	$0.73 \pm 0.10^{+0.15}_{-0.05}$
Rel. phase,	$53 \pm 61^{+5}_{-50}$	$-20 \pm 18^{+14}_{-9}$	$6 \pm 24^{+23}_{-59}$

Resonant substructure $h_b(\text{nP})\pi^\pm\pi^\mp$





probe: missing mass

$$e^+ e^- \rightarrow \underbrace{h_b(\text{nP})\pi^\pm\pi^\mp}_{\text{"Z"}}$$

$$M_Z = MM(\pi) = \sqrt{E_Z^2 - p_Z^2}$$

yield in $MM(\pi)$ bins




Fit function: $|BW(s, M_1, \Gamma_1) + ae^{i\phi} BW(s, M_2, \Gamma_2) + be^{i\psi}|^2 \frac{qp}{\sqrt{s}}$

Nearly all resonant

2 new resonances: $Z_b^\pm(10610)$, $Z_b^\pm(10650)$, 5 modes ea




21



Good
agreement
among all 5
modes

- Relative phases: Υ ($\sim 0^\circ$), h_b ($\sim 180^\circ$)
- Masses just above B^*B and B^*B^* thresholds
- angular analysis favors $J^P=1^+$ [arXiv:1403.0992]
 Favors "meson molecule" hypothesis of Z_b 's

Further evidence: neutral partner



$$e^+ e^- \rightarrow \Upsilon(nS) \pi^0 \pi^0$$

$$M(\Upsilon(2S)\pi^0)$$

$$Z_b^0 \rightarrow \Upsilon(nS) \pi^0$$

$Z_b^0(10610)$ observed with 6.5σ significance



[PRD 88, 052016 (2013)]

Further evidence $Z_b \rightarrow B^* B^{(*)}$



$e^+ e^- \rightarrow B^* B^{(*)} \pi^\pm$

121.4 fb^{-1}



arXiv:1209.6450 [hep-ex]

What IS Z_b ?

many theories

- Meson molecule
[A.Bondar, et al., PRD 84, 054010 (2011)]
- Coupled channel resonances
[I.V.Danilkin et al, arXiv:1106.1552]
- Cusp
[D.Bugg, Europhys.Lett. 96, 11002 (2011)]
- Tetraquark
[M.Karliner & H.Lipkin, arXiv:0802.0649]

Similar pattern in charmonium region

Heavy quark exotica

PDG13

Many are unconfirmed
Primary characteristic:
high rate to quarkonia

Charmonium
-like

Z(3900)
Z(3885)
Z(4025)
Z(4020)

Bottomonium-
like

Z_b⁰(10610)

State	m (MeV)	Γ (MeV)	J^{PC}	Process (mode)	Experiment (# σ)	Year	Status
$X(3872)$	3871.68 ± 0.17	< 1.2	$1^{++}/2^{-+}$	$B \rightarrow K(\pi^+\pi^- J/\psi)$ $p\bar{p} \rightarrow (\pi^+\pi^- J/\psi) + \dots$ $B \rightarrow K(\omega J/\psi)$ $B \rightarrow K(D^{*0}\bar{D}^0)$ $B \rightarrow K(\gamma J/\psi)$ $B \rightarrow K(\gamma\psi(2S))$ $pp \rightarrow (\pi^+\pi^- J/\psi) + \dots$	Belle [36,37] (12.8), BABAR [38] (8.6) CDF [39–41] (np), D0 [42] (5.2) Belle [43] (4.3), BABAR [23] (4.0) Belle [44,45] (6.4), BABAR [46] (4.9) Belle [47] (4.0), BABAR [48,49] (3.6) BABAR [49] (3.5), Belle [47] (0.4) LHCb [50] (np)	2003	OK
$X(3915)$	3917.4 ± 2.7	28_{-9}^{+10}	$0/2^{?+}$	$B \rightarrow K(\omega J/\psi)$ $e^+e^- \rightarrow e^+e^-(\omega J/\psi)$	Belle [51] (8.1), BABAR [52] (19)	2004	OK
$X(3940)$	3942_{-8}^{+9}	37_{-17}^{+27}	$?^{?+}$	$e^+e^- \rightarrow J/\psi(D\bar{D}^*)$ $e^+e^- \rightarrow J/\psi(\dots)$	Belle [53] (7.7), BABAR [23] (np) Belle [54] (6.0) Belle [20] (5.0)	2007	NC!
$G(3900)$	3943 ± 21	52 ± 11	1^{--}	$e^+e^- \rightarrow \gamma(D\bar{D})$	BABAR [55] (np), Belle [56] (np)	2007	OK
$Y(4008)$	4008_{-49}^{+121}	226 ± 97	1^{--}	$e^+e^- \rightarrow \gamma(\pi^+\pi^- J/\psi)$	Belle [57] (7.4)	2007	NC!
$Z_1(4050)^+$	4051_{-43}^{+24}	82_{-55}^{+51}	$?$	$B \rightarrow K(\pi^+\chi_{c1}(1P))$	Belle [58] (5.0), BABAR [59] (1.1)	2008	NC!
$Y(4140)$	4143.4 ± 3.0	15_{-7}^{+11}	$?^{?+}$	$B \rightarrow K(\phi J/\psi)$	CDF [60,61] (5.0)	2009	NC!
$X(4160)$	4156_{-25}^{+29}	139_{-65}^{+113}	$?^{?+}$	$e^+e^- \rightarrow J/\psi(D\bar{D}^*)$	Belle [54] (5.5)	2007	NC!
$Z_2(4250)^+$	4248_{-45}^{+185}	177_{-72}^{+321}	$?$	$B \rightarrow K(\pi^+\chi_{c1}(1P))$	Belle [58] (5.0), BABAR [59] (2.0)	2008	NC!
$Y(4260)$	4263_{-9}^{+8}	95 ± 14	1^{--}	$e^+e^- \rightarrow \gamma(\pi^+\pi^- J/\psi)$	BABAR [62,63] (8.0)	2005	OK
				$e^+e^- \rightarrow (\pi^+\pi^- J/\psi)$ $e^+e^- \rightarrow (\pi^0\pi^0 J/\psi)$	CLEO [64] (5.4), Belle [57] (15)		
$Y(4274)$	$4274.4_{-6.7}^{+8.4}$	32_{-15}^{+22}	$?^{?+}$	$B \rightarrow K(\phi J/\psi)$	CLEO [65] (11)		
$X(4350)$	$4350.6_{-5.1}^{+4.6}$	$13.3_{-10.0}^{+18.4}$	$0/2^{++}$	$e^+e^- \rightarrow e^+e^-(\phi J/\psi)$	CDF [61] (3.1)	2010	NC!
$Y(4360)$	4361 ± 13	74 ± 18	1^{--}	$e^+e^- \rightarrow \gamma(\pi^+\pi^-\psi(2S))$	Belle [66] (3.2)	2009	NC!
$Z(4430)^+$	4443_{-18}^{+24}	107_{-71}^{+113}	$?$	$B \rightarrow K(\pi^+\psi(2S))$	BABAR [67] (np), Belle [68] (8.0)	2007	OK
$X(4630)$	4634_{-11}^{+9}	92_{-32}^{+41}	1^{--}	$e^+e^- \rightarrow \gamma(\Lambda_c^+\Lambda_c^-)$	Belle [69,70] (6.4), BABAR [71] (2.4)	2007	NC!
$Y(4660)$	4664 ± 12	48 ± 15	1^{--}	$e^+e^- \rightarrow \gamma(\pi^+\pi^-\psi(2S))$	Belle [72] (8.2)	2007	NC!
$Z_b(10610)^+$	10607.2 ± 2.0	18.4 ± 2.4	1^+	$\Upsilon(5S) \rightarrow \pi^-(\pi^+[b\bar{b}])$	Belle [68] (5.8)	2007	NC!
$Z_b(10650)^+$	10652.2 ± 1.5	11.5 ± 2.2	1^+	$\Upsilon(5S) \rightarrow \pi^-(\pi^+[b\bar{b}])$	Belle [73,74] (16)	2011	NC!
$Y_b(10888)$	10888.4 ± 3.0	$30.7_{-7.7}^{+8.9}$	1^{--}	$e^+e^- \rightarrow (\pi^+\pi^-\Upsilon(nS))$	Belle [73,74] (16)	2011	NC!
					Belle [75,76] (2.0)	2010	NC!



Anomalies of the $\Upsilon(10860)$

energy scan


$\sigma_{b\bar{b}} \text{ vs } \sigma_{\Upsilon\pi\pi}$

[PRD82, 091106 (2010)]

$$\sigma(b\bar{b})$$

Event count

$$(3.40 \pm 0.16) \times 10^5 \text{ events/fb}^{-1} \quad (121.4 \text{ fb}^{-1})$$



Event shape parameter
(Fox-Wolfram moments)

$$R_2 = \frac{\sum_{i,j} |p_i||p_j| P_2(\cos \theta)}{\sum_{i,j} |p_i||p_j| P_0(\cos \theta)}$$

2-jet $e^+e^- \rightarrow q\bar{q}$ $R_2 \rightarrow 1$

$e^+e^- \rightarrow B\bar{B}$ $R_2 \rightarrow 0$

Contributions of initial-state radiation calculated, subtracted

$e^+e^- \rightarrow \gamma\gamma^*$


$\gamma^* \rightarrow \Upsilon(1/2/3S)$

scan near $\Upsilon(10860)$ [PRD82, 091106 (2010)]

$\sqrt{s}(\text{GeV})$	$\mathcal{L}(\text{fb}^{-1})$
10.8275	1.68
10.8825	1.83
10.8975	1.41
10.9275	1.14
10.9575	1.01
11.0175	0.86

$$M_{5S}c^2 = [10888^{+2.7}_{-2.6} \pm 1.2] \text{ MeV}$$

$$\Delta Mc^2 = 9 \pm 4 \text{ MeV}$$





Anomalies of the $\Upsilon(10860)$

energy scan

$$\sigma_{b\bar{b}} \text{ vs } \sigma_{\Upsilon\pi\pi}$$

-> reprise 121.4 fb^{-1} @ $10.865 \pm 1 \text{ GeV}$
 $+15 \times 1 \text{ fb}^{-1}, +61 \times 50 \text{ pb}^{-1}$ @ $10.68-10.11.02$


arXiv:1501.01137



$$R_b \equiv \frac{\sigma(e^+e^- \rightarrow b\bar{b})}{\sigma(e^+e^- \rightarrow \mu^+\mu^-)}$$

Fitting Model:
 2 Breit-Wigner+
 Flat continuum
 (coherent+incoherent)
 [match Babar]

R_b



$$|A_{NR}|^2 +$$

$$|A_R + A_{5S} e^{i\phi_{5S}} BW(M_{5S}, \Gamma_{5S}) + A_{6S} e^{i\phi_{6S}} BW(M_{6S}, \Gamma_{6S})|^2$$




 $\sigma(b\bar{b})$

$$R_b \equiv \frac{\sigma(e^+e^- \rightarrow b\bar{b})}{\sigma(e^+e^- \rightarrow \mu^+\mu^-)}$$

$$M_{5S} = 10881.8^{+1.0}_{-1.1} \pm 1.2$$

$$\Gamma_{5S} = 48.5^{+1.9+2.0}_{-1.8-2.8}$$

$$M_{6S} = 11003.0 \pm 1.1^{+0.9}_{-1.0}$$

$$\Gamma_{6S} = 39.3^{+1.7+1.3}_{-1.6-2.4}$$

R_b
0.5
0.4
0.2
0
10.7
10.75
10.8
10.85
10.9
10.95
11
11.05
Υ(10860)
Υ(11020)
NR
R
Cross term w A_R
Υ(10860)
Υ(11020)
√s (GeV)


$$|A_{NR}|^2 +$$

$$|A_R + A_{5S}e^{i\phi_{5S}}BW(M_{5S}, \Gamma_{5S}) + A_{6S}e^{i\phi_{6S}}BW(M_{6S}, \Gamma_{6S})|^2$$

$e^+e^- \rightarrow \Upsilon(nS)\pi^\pm\pi^\mp$



$\Upsilon(nS) \rightarrow \mu^+\mu^-$



$$e^+ e^- \rightarrow \Upsilon(nS) \pi^\pm \pi^\mp$$

- event-by-event efficiency correction over Dalitz space,
(reduce model-dependence)

- fit (simultaneous for 3 Υ 's)

$$\underline{PHSP}(E_{CM}) \times (|A_{NR}|^2$$

$$+ |A_R + A_{5S} e^{i\phi_{5S}} BW(M_{5S}, \Gamma_{5S}) + A_{6S} e^{i\phi_{6S}} BW(M_{6S}, \Gamma_{6S})|^2)$$

- set $|A_{NR}|, |A_R| = 0$

- possible differences in Dalitz distribution btw
 $\Upsilon(10860), \Upsilon(11020)$
-> "decoherence coefficient" = $k e^{i\delta}$ ($0 < k < 1$)

$$PHSP(E_{CM}) \times |A_{5S} BW_{5S}|^2 + |A_{6S} BW_{6S}|^2$$

$$+ 2k A_{5S} A_{6S} \Re[e^{i\delta} BW_{5S} BW_{6S}^*]$$

- (k consistent w 1)

Masses, widths



$$R_{\Upsilon\pi\pi} \equiv \frac{\sigma(e^+e^- \rightarrow \Upsilon(nS)\pi^+\pi^-)}{\sigma(e^+e^- \rightarrow \mu\mu)}$$

$$M_{5S} = 10891.1 \pm 3.2^{+0.6}_{-1.5}$$


$$\Gamma_{5S} = 53.7^{+7.1+0.9}_{-5.6-5.4}$$

$$M_{6S} = 10987.5^{+6.4+9.0}_{-2.5-2.1}$$

$$\Gamma_{6S} = 61^{+9+2}_{-19-20}$$

$\Upsilon\pi\pi$ vs $b\bar{b}$

$$\Delta M c^2 = 9.3 \pm 3.9 \text{ MeV}$$





Anomalies of the $\Upsilon(10860)$

High rates to

$$\Upsilon(nS)\pi\pi$$

$$h_b(mP)\pi\pi$$

$$B^* B^{(*)}\pi$$

Large fraction as $Z_b\pi$ (X in PDG as of 2014)
energy scan:

$\Upsilon(10860), \Upsilon(11020)$ compatible masses, widths

$\sigma_{b\bar{b}}$ large continuum

$\sigma_{\Upsilon\pi\pi} \sim$ no continuum

-> Reconcile resonance amplitudes

Contribution of bottomonium modes to $\Upsilon(10860)$ resonance


$$PHSP(E_{CM}) \times |A_{5S}BW_{5S}|^2 + |A_{6S}BW_{6S}|^2 + 2kA_{5S}A_{6S}\Re[e^{i\delta}BW_{5S}BW_{6S}^*]$$

$$\mathcal{P} \equiv \frac{\sum_n P_n}{P_b} = 0.42 \pm 0.04 \quad \text{all } \Upsilon\pi\pi, h_b\pi\pi$$

$$= 1.09 \pm 0.15 \quad \text{with } B^*B^{(*)}\pi$$

??Appears to saturate b cross section
w/o $B^{(*)}, B_s^{(*)}$ pairs

Look again at fit R_b




Paradox?

large interference,
cannot come from
bottomonium modes

but final states w
bottomonium/
Bottomonium-like
saturate
amplitude?

Look again at fit R_b



Paradox?

Model of
2 resonances + flat
continuum
Cannot be used to
describe this
region,
Despite great fit,
consistent M, Γ

$$|A_{NR}|^2 +$$

$$|A_R + A_{5S}e^{i\phi_{5S}}BW(M_{5S}, \Gamma_{5S}) + A_{6S}e^{i\phi_{6S}}BW(M_{6S}, \Gamma_{6S})|^2$$

Summary

39

Rich structure in region of $\Upsilon(10860)$

- $e^+e^- \rightarrow \{b\bar{b}\}\pi\pi$

New states $Z_b^+(10610), Z_b^+(10650)$

Seen to decay to BB^*, B^*B^*

Evidence for neutral Z_b

- new questions raised by $R_b, R_{\Upsilon\pi\pi}$ vs E_{CM}
what is $\Upsilon(10860)$?
 - High rate to bottomonia, Z_b 's $\rightarrow Y_b$?
 - masses from $R_b, R_{\Upsilon\pi\pi}$ consistent
but rates are a puzzle
 - use $R_{\Upsilon\pi\pi}$ as primary measurement of M, Γ
- to be continued...
 $B^{(*)}_{(s)}$ modes vs E_{CM}



Backup Slides